

Pacific bluefin tuna transport Fukushima-derived radionuclides from Japan to California

Daniel J. Madigan^{a,1}, Zofia Baumann^b, and Nicholas S. Fisher^b

^aHopkins Marine Station, Stanford University, Pacific Grove, CA 93950; and ^bSchool of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11794

Edited by Karl K. Turekian, Yale University, North Haven, CT, and approved April 25, 2012 (received for review March 22, 2012)

The Fukushima Dai-ichi release of radionuclides into ocean waters caused significant local and global concern regarding the spread of radioactive material. We report unequivocal evidence that Pacific bluefin tuna, *Thunnus orientalis*, transported Fukushima-derived radionuclides across the entire North Pacific Ocean. We measured γ -emitting radionuclides in California-caught tunas and found ^{134}Cs ($4.0 \pm 1.4 \text{ Bq kg}^{-1}$) and elevated ^{137}Cs ($6.3 \pm 1.5 \text{ Bq kg}^{-1}$) in 15 Pacific bluefin tuna sampled in August 2011. We found no ^{134}Cs and background concentrations ($\sim 1 \text{ Bq kg}^{-1}$) of ^{137}Cs in pre-Fukushima bluefin and post-Fukushima yellowfin tunas, ruling out elevated radiocesium uptake before 2011 or in California waters post-Fukushima. These findings indicate that Pacific bluefin tuna can rapidly transport radionuclides from a point source in Japan to distant ecoregions and demonstrate the importance of migratory animals as transport vectors of radionuclides. Other large, highly migratory marine animals make extensive use of waters around Japan, and these animals may also be transport vectors of Fukushima-derived radionuclides to distant regions of the North and South Pacific Oceans. These results reveal tools to trace migration origin (using the presence of ^{134}Cs) and potentially migration timing (using ^{134}Cs : ^{137}Cs ratios) in highly migratory marine species in the Pacific Ocean.

pelagic | radioactivity | bioaccumulation

The infrequency of nuclear accidents coupled with potentially wide-ranging effects on ecosystems and human health make the dynamics and risks of radionuclide discharge into the environment a relatively poorly understood but highly important area of research (1–4). On March 11, 2011, an earthquake and subsequent tsunami led to flooding of the Fukushima Dai-ichi nuclear power plants in Japan. Coolant pumps failed to operate and the power plant reactors overheated, leading to a release of radionuclides directly into the ocean exceeding that from any previous accident (3). The release of radionuclides produced a 1–2 wk pulse that peaked on April 6, 2011 with ocean concentrations of 68 MBq m^{-3} (3) and an estimated total release of up to $22 \times 10^{15} \text{ Bq}$ of ^{137}Cs (4) ($1 \text{ Bq} = 1 \text{ disintegration s}^{-1}$). The dominant long-lived γ -emitting radionuclides $^{134}\text{Cesium}$ ($t_{1/2} = 2.1 \text{ y}$) and ^{137}Cs ($t_{1/2} = 30 \text{ y}$) were released at a consistent ratio of about 1 (0.99 ± 0.03) (3), although the ratio of ^{134}Cs : ^{137}Cs would decline slowly over time because of the differential decay rate constants of the two isotopes (Fig. S1). After considerable dilution 2–3 mo after maximum discharge, surface concentrations still exceeded prior concentrations by up to 10,000-fold in coastal waters (3) and up to 1,000-fold over a $150,000 \text{ km}^2$ area of the Pacific up to 600 km east of Japan (4). Before the Fukushima discharge, low concentrations (1.5 mBq L^{-1}) of the long-lived ^{137}Cs (fallout from weapons testing) were detectable in Japanese waters (3), whereas the shorter-lived ^{134}Cs was undetectable in Pacific surface waters and biota.

The Pacific bluefin tuna (PBFT), *Thunnus orientalis*, is a highly migratory fish that inhabits the western and eastern North Pacific Ocean at various life stages (5) (Fig. 1A). Mature PBFT spawn in the western Pacific, and some juveniles remain in Japanese waters while others migrate eastward to the California Current Large Marine Ecosystem (CCLME) (Fig. 1A), with most migrating late

in their first year or early in their second (5). Thus, all bluefin between years 1–2 (here, 2-y-old PBFT) caught during summer in the eastern Pacific must have migrated from the western Pacific within several months of capture. Waters north of the Kuroshio Current (Fig. 1A) showed high radionuclide concentrations in spring 2011 (3), and juveniles make extensive use of this region before their eastward migration to the CCLME (6).

We tested the possibility that juvenile PBFT served as biological vectors of radionuclides between two distant ecoregions: the waters off Japan and the CCLME. We analyzed 2-y-old PBFT caught off San Diego, CA, in August 2011, known from size to be recent Japan migrants, for the presence of Fukushima-derived radionuclides. Because Cs accumulates in the muscle tissue of fish (7), we analyzed the white muscle tissue of PBFT in 2011 for concentrations of ^{134}Cs , ^{137}Cs , and various naturally occurring γ -emitting radionuclides. To rule out non-Fukushima sources of radiocesium in fish muscle, we also measured radionuclide concentrations in PBFT collected in California waters before the Fukushima discharge (2008) and in yellowfin tuna (YFT), *T. albacares* (August 2011), in the CCLME where they are highly residential (8, 9).

Results and Discussion

All 15 PBFT collected in 2011 contained ^{134}Cs ($4.0 \pm 1.4 \text{ Bq kg}^{-1}$ dry wt) and ^{137}Cs ($6.3 \pm 1.5 \text{ Bq kg}^{-1}$) in white muscle tissue (Table 1, Table S1, and Fig. 1A). At the time of capture, total $^{134}+^{137}\text{Cs}$ concentrations were about 10 times higher in 2011 PBFT than in PBFT from previous years (Table 1). In contrast, 2008 PBFT and 2011 YFT had no measurable ^{134}Cs and consistent, much lower ^{137}Cs concentrations (consistent with background concentrations from fallout) than the 2011 PBFT (Table 1, Table S2, and Fig. 1A). This is unequivocal evidence that Fukushima-derived radionuclides were transported to the CCLME by Pacific bluefin tuna, as no other sources of ^{134}Cs were present in the North Pacific preceding the Fukushima disaster (3, 4). Mean concentrations of the naturally occurring γ -emitting ^{40}K in the 2011 PBFT were $347 \pm 49 \text{ Bq kg}^{-1}$ (Table 1). Other naturally occurring γ -emitting radionuclides (^7Be , ^{211}Bi , and ^{212}Pb) were detectable at extremely low concentrations (Table S3), approximately three orders of magnitude below measured radiocesium concentrations.

Because bluefin tuna are harvested annually in the Eastern Pacific Ocean (EPO) at $1.7\text{--}9.9 \times 10^3$ metric tonnes (10) (Table S4) for human consumption (2000 to 2010), the possibility of radioactive contamination raises public health concerns. Radiocesium concentrations of post-Fukushima PBFT reported here were more than an order of magnitude below the recently

Author contributions: D.J.M. and N.S.F. designed research; D.J.M., Z.B., and N.S.F. performed research; N.S.F. contributed new reagents/analytic tools; D.J.M., Z.B., and N.S.F. analyzed data; and D.J.M., Z.B., and N.S.F. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: dmadigan@stanford.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1204859109/-DCSupplemental.

Table 1. Measured ^{134}Cs , ^{137}Cs , and the naturally occurring radionuclide ^{40}K for post-Fukushima bluefin (PBFT 2011), pre-Fukushima bluefin (PBFT 2008), and post-Fukushima yellowfin tuna (YFT 2011) caught in California waters

	SL (cm)*	Body mass (kg dry) [†]	Age (y) [‡]	Radionuclide concentrations (Bq kg ⁻¹)				
				^{134}Cs	^{137}Cs	^{40}K	$^{134}\text{Cs}:^{137}\text{Cs}$	$^{134+137}\text{Cs}$
PBFT 2011 (n = 15)								
Median	66.5	1.5	1.5	4.3	6.0	367	0.66	10.6
Mean	66.2	1.5	1.5	4.0	6.3	347	0.62	10.3
SD	3.6	0.2	0.1	1.4	1.5	49	0.14	2.9
PBFT 2008 (n = 5)								
Median	66.3	1.5	1.4	0	1.4	266	0	1.37
Mean	66.2	1.5	1.4	0	1.4	258	0	1.11
SD	1.2	0.09	0.05	0	0.2	43	0	0.64
YFT 2011 (n = 5)								
Median	72.3	1.9	1.2	0	1.2	342	0	1.21
Mean	72.3	1.9	1.2	0	1.1	333	0	1.10
SD	2.5	0.2	0.01	0	0.4	78	0	0.35

All radionuclide concentrations are in Bq kg⁻¹ dry mass.
 *Estimated from CFL [PBFT (20) and YFT (21)].
[†]Estimated from standard length (SL) (22, 23).
[‡]Estimated from SL (24, 25).

model performs well in predicting previous concentrations of Cs in tuna captured months after exposure.

Decay-corrected $^{134}\text{Cs}:^{137}\text{Cs}$ ratios measured in post-Fukushima bluefin caught off California averaged 0.62 ± 0.14 and ranged from 0.26 to 0.84 (Table 1 and Table S1). Ratios of $^{134}\text{Cs}:^{137}\text{Cs}$ have been used to discern sedimentation patterns and water movements in the Irish Sea, which received 15 PBq of discharged ^{137}Cs between 1978 and 1983 from the nuclear fuel-reprocessing plant at Sellafield (15, 16). However, Cs ratios have not been used previously to track biota movements. Assuming that PBFT tissue would reflect the $^{134}\text{Cs}:^{137}\text{Cs}$ ratio of 1.0 to which they were exposed near Japan, radiocesium ratios can be used to make inferences about migration and the timing of exposure to contaminated waters. Back-calculated $^{134}\text{Cs}:^{137}\text{Cs}$ ratios in PBFT

approached 1.0 (0.8 ± 0.2) at 4 mo before capture (Fig. 2), suggesting departure from Japan ~4 mo before capture off California. This result suggests that the radiocesium levels in California-caught PBFT were the result of <1 mo exposure to contaminated waters near Japan. Because radiocesium release was pulsed, it is possible that back-calculated ratios will always converge on the period of maximum discharge (rather than on the date of PBFT emigration from Japanese waters). The $^{134}\text{Cs}:^{137}\text{Cs}$ ratio of seawater near Japan would have changed little from 1.0 within 4 mo of the pulsed radioactivity discharge (1.0–0.9; Fig. S1). However, additional PBFT radiocesium data across longer time scales (e.g., PBFT from 2012) will indicate whether back-calculated $^{134}\text{Cs}:^{137}\text{Cs}$ ratios in PBFT consistently converge on 1.0 in April 2011 (period of maximum discharge) or converge on 1.0




Fig. 2. Measured (corrected for background levels of ^{137}Cs) and back-calculated estimates of radiocesium concentrations in muscle of post-Fukushima Pacific bluefin tuna *Thunnus orientalis* (n = 15). Estimated values of ^{134}Cs and ^{137}Cs concentrations in 2011 PBFT for various times before capture in California (30, 60, 90, 120 d) account for background levels of ^{137}Cs , decreases in tissue concentrations because of growth dilution and efflux rates of radiocesium out of the fish during their migration across the Pacific (detailed in SI Methods). Mean concentrations of ^{134}Cs (red triangles) or ^{137}Cs (gray squares) shown on the left y axis. Error bars represent 1 SD. Ratios of $^{134}\text{Cs}:^{137}\text{Cs}$ for each individual fish (empty circles) shown with scale on the right y axis. Dotted line represents 1:1 ratio of $^{134}\text{Cs}:^{137}\text{Cs}$, the ratio expected in tuna while in waters off Japan contaminated with radiocesium at a $^{134}\text{Cs}:^{137}\text{Cs}$ ratio of 1.0.

over a range of dates after the Fukushima discharge (demonstrating the utility of ^{134}Cs : ^{137}Cs ratios as a tool to assess the timing of PBFT migrations).

The total load of radiocesium transported to the CCLME by PBFT can be estimated from catches in the EPO. Catch data varies yearly (10), but assuming PBFT commercial catches in 2011 were within the range of catch from 2000 to 2010, transported and harvested radiocesium in tuna muscle tissue in 2011 could range from $\sim 3\text{--}17 \times 10^6$ Bq (Table S4) or $\ll 1\%$ of total radiocesium released into Japanese waters (4). Catch data represent a portion of the PBFT in the EPO, so total transport of radiocesium by PBFT would likely be higher. Still, this is a small quantity of radiocesium to be introduced to a large pelagic ecosystem, but it is also a conservative estimate based on one species. Other highly migratory species (HMS) (e.g., turtles, sharks, and seabirds) that forage near Japan may assimilate radiocesium and transport it to distant regions of the north and south Pacific (Fig. 1B). Tissue concentrations of radiocesium in these species would depend on time spent near Japan, foraging strategies, and timing of migration. The potential for species in Fig. 1B and other HMS (e.g., pinnipeds, whales, and billfish) that forage in Japan to transport Fukushima-derived radiocesium is speculative. However, the presence of Fukushima-derived radiocesium in all 2011 PBFT individuals reported here suggests that study of other HMS is warranted. Our results demonstrate that Fukushima-derived radionuclides in animal tissues can serve as tracers of both migration origin (presence or absence of ^{134}Cs) and, potentially, timing (using ^{134}Cs : ^{137}Cs ratios) in mobile marine animals, providing valuable complementary movement data to extensive tagging programs in the Pacific (8). Extensive data regarding spatiotemporal variations in Cs concentrations in the west Pacific, and consequent uptake by biota, are forthcoming, which will sharpen the precision of these tracers. The Fukushima

disaster, thus, provides an opportunity to examine both the extent of transport of anthropogenic radionuclides by highly migratory species and an unexpected tool for examining migratory origins of apex predators in the Pacific Ocean.

Methods

Tuna tissue samples were collected from recreational anglers in San Diego, CA. Muscle samples were freeze-dried and ground with mortar and pestle and analyzed using a low-energy germanium detector. We detected the γ emissions of ^{137}Cs and ^{134}Cs , ^{40}K , and other naturally occurring radionuclides. ^{134}Cs and ^{137}Cs concentrations of post-Fukushima PBFT and YFT samples were decay-corrected to the angler-estimated catch date for all fish. We calculated ^{134}Cs and ^{137}Cs concentrations in 2011 PBFT for various times before capture in California (0, 30, 60, 90, 120 d). We accounted for background ^{137}Cs in tuna muscle (1.0 Bq kg^{-1}) by subtracting 1.0 from total ^{137}Cs values. We then accounted for the radioactive decay of Cs isotopes, important only for the shorter-lived ^{134}Cs ($t_{1/2} = 2.1 \text{ y}$) using an exponential decay model. To account for the metabolic efflux of assimilated cesium out of fish, we used an experimentally derived marine fish efflux rate constant k of 0.019 d^{-1} (14). To address growth dilution of the Cs concentrations in muscle, we calculated change in fish body mass (0–120 d before catch) and calculated dilution of Cs concentrations over this range of days due to growth.

We estimated radiocesium transport to the CCLME by PBFT using catch data as a proxy for PBFT biomass in the CCLME. Catch data from 2000 to 2010 were converted to muscle biomass and then multiplied by mean measured radiocesium concentrations to generate a range of estimates of transported radiocesium. Spearman's ρ analyses (two-tailed, nonparametric; $\alpha = 0.05$) were performed to assess correlation between 2011 PBFT length and radionuclide concentrations.

Equations and further details of the methods are included in *SI Methods*.

ACKNOWLEDGMENTS. We thank S. Zegers for laboratory assistance; A. Carlisle for technical assistance; and K. Buesseler, M. Condrup, H. Dewar, F. Micheli, and two anonymous reviewers for helpful comments. We also thank the anonymous fishermen who donated fish samples toward this study. This work was funded by the Gordon and Betty Moore Foundation.

- Garnier-Laplace J, Beaugelin-Seiller K, Hinton TG (2011) Fukushima wildlife dose reconstruction signals ecological consequences. *Environ Sci Technol* 45:5077–5078.
- Masson O, et al. (2011) Tracking of airborne radionuclides from the damaged Fukushima Dai-ichi nuclear reactors by European networks. *Environ Sci Technol* 45:7670–7677.
- Buesseler K, Aoyama M, Fukasawa M (2011) Impacts of the Fukushima nuclear power plants on marine radioactivity. *Environ Sci Technol* 45:9931–9935.
- Buesseler KO, et al. (2012) Fukushima-derived radionuclides in the ocean and biota off Japan. *Proc Natl Acad Sci USA* 109:5984–5988.
- Bayliff WH (1994) A review of the biology and fisheries for northern bluefin tuna, *Thunnus thynnus*, in the Pacific Ocean. *FAO Fish Tech Pap* 336(2):244–295.
- Kitagawa T, et al. (2009) Immature Pacific bluefin tuna, *Thunnus orientalis*, utilizes cold waters in the Subarctic Frontal Zone for trans-Pacific migration. *Environ Biol Fishes* 84:193–196.
- Young DR, Folsom TR, Hodge VF (1975) ^{137}Cs and ^{40}K in the flesh of Pacific albacore, 1964–1974. *Health Phys* 29:689–694.
- Block BA, et al. (2011) Tracking apex marine predator movements in a dynamic ocean. *Nature* 475:86–90.
- Schaefer KM, Fuller DW, Block BA (2007) Movements, behavior, and habitat utilization of yellowfin tuna (*Thunnus albacares*) in the northeastern Pacific Ocean, ascertained through archival tag data. *Mar Biol* 152:503–525.
- Inter-American Tropical Tuna Commission (2010) *Fishery Status Report* (Inter-American Tropical Tuna Commission, La Jolla, CA).
- Japanese Ministry of Agriculture, Forestry, and Fisheries (2011) Results of the inspection on radioactivity materials in fisheries products. Available at <http://www.jfa.maff.go.jp/e/inspection/index.html>.
- Aarkrog A, et al. (1997) A comparison of doses from ^{137}Cs and ^{210}Po in marine food: A major international study. *J Environ Radioact* 34:69–90.
- Stewart GM, Fowler SW, Fisher NS (2008) *U-Th Series Nuclides in Aquatic Systems*, eds Krishnaswami S, Cochran JK (Elsevier, Amsterdam), pp 269–305.
- Mathews T, Fisher NS (2009) Dominance of dietary intake of metals in marine elasmobranch and teleost fish. *Sci Total Environ* 407:5156–5161.
- Aston SR, Stanners DA (1979) The determination of estuarine sedimentation rates by ^{134}Cs / ^{137}Cs and other artificial radionuclide profiles. *Estuar Coast Mar Sci* 9:529–541.
- Smith JN, Ellis KM, Jones EP (1990) Cesium 137 transport into the Arctic Ocean through Fram Strait. *J Geophys Res* 95:1693–1701.
- Nagasawa K (1998) Predation by salmon sharks (*Lamna ditropis*) on Pacific salmon (*Oncorhynchus* spp.) in the North Pacific Ocean. *North Pac Anadr Fish Comm Bull* 1:419–433.
- Shaffer SA, et al. (2006) Migratory shearwaters integrate oceanic resources across the Pacific Ocean in an endless summer. *Proc Natl Acad Sci USA* 103:12799–12802.
- Peckham SH, et al. (2007) Small-scale fisheries bycatch jeopardizes endangered Pacific loggerhead turtles. *PLoS ONE* 2:e1041.
- Farwell CJ (2011) *Based on a Regression by C. Farwell* (Tuna Research and Conservation Center, Pacific Grove, CA).
- Scida P, Rainosek A, Lowery T (2001) Length conversions for yellowfin tuna (*Thunnus albacares*) caught in the western North Atlantic Ocean. *Collect Vol Sci Pap ICCAT* 52:528–532.
- Itoh T (2001) Estimation of total catch in weight and catch-at-age in number of bluefin tuna *Thunnus orientalis* in the whole Pacific Ocean. *Bull Natl Res Inst Far Seas Fish* 38:83–111.
- Wild A (1986) Growth of yellowfin tuna, *Thunnus albacares*, in the Eastern Pacific Ocean based on otolith increments. *Inter-Am Trop Tuna Comm Bull* 18:421–482.
- Bayliff WH, Ishizuka Y, Deriso R (1991) Growth, movement, and attrition of northern bluefin tuna, *Thunnus thynnus*, in the Pacific ocean, as determined by tagging. *Inter-Am Trop Tuna Comm Bull* 20:3–94.
- Wild A (1994) A review of the biology and fisheries for yellowfin tuna, *Thunnus albacares*, in the Eastern Pacific Ocean. *FAO Fish Tech Pap* 336:52–107.